antibodies.com

Datasheet for ABIN7176223 anti-ZNF3 antibody (AA 92-210)

Image



Overview

Quantity:	100 µg
Target:	ZNF3
Binding Specificity:	AA 92-210
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF3 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Zinc finger protein 3 protein (92-210AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details


Target:	ZNF3
Alternative Name:	ZNF3 (ZNF3 Products)
Background:	Background: Involved in cell differentiation and/or proliferation.
	Aliases: ZNF3 antibody, KOX25 antibody, Zinc finger protein 3 antibody, Zinc finger protein

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7176223 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details

	HF.12 antibody, Zinc finger protein HZF3.1 antibody, Zinc finger protein KOX25 antibody
UniProt:	P17036
Application Details	
Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7176223 diluted at 1:300 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.