

Datasheet for ABIN7176263
anti-ZNF367 antibody (AA 83-151)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	ZNF367
Binding Specificity:	AA 83-151
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF367 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Zinc finger protein 367 protein (83-151AA)
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	ZNF367
Alternative Name:	ZNF367 (ZNF367 Products)
Background:	Background: Transcriptional activator. Isoform 1 may be involved in transcriptional activation of erythroid genes.

Target Details

Aliases: C2H2 zinc finger protein ZFF29 antibody, Zinc finger protein 367 antibody, ZN367_HUMAN antibody, znf367 antibody

UniProt: [Q7RTV3](#)

Application Details

Application Notes: Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7176263 diluted at 1:400 and staining in paraffin-embedded human glioma performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blotting

Image 2. Western Blot Positive WB detected in: K562 whole cell lysate, NIH/3T3 whole cell lysate All lanes: ZNF367 antibody at 6 µg/mL Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 39, 33 kDa Observed band size: 39 kDa