antibodies .- online.com

Datasheet for ABIN7176620 anti-ZKSCAN5 antibody (AA 490-643)

2 Images



Overview

Quantity:	100 µg
Target:	ZKSCAN5
Binding Specificity:	AA 490-643
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZKSCAN5 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Recombinant Human Zinc finger protein with KRAB and SCAN domains 5 protein (490-643AA)
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details


Target:	ZKSCAN5
Alternative Name:	ZKSCAN5 (ZKSCAN5 Products)
Background:	Background: May be involved in transcriptional regulation.
	Aliases: ZKSCAN5 antibody, KIAA1015 antibody, ZFP95 antibody, Zinc finger protein with KRAB

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7176620 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details


	and SCAN domains 5 antibody, Zinc finger protein 95 homolog antibody, Zfp-95 antibody
UniProt:	Q9Y2L8
Application Details	
Application Notes:	Recommended dilution: IHC:1:200-1:500,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Preservative: 0.03 % Proclin 300
	Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunohistochemistry

Image 1. IHC image of ABIN7176620 diluted at 1:200 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry

Image 2. IHC image of ABIN7176620 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.