

Datasheet for ABIN7176651
anti-ZFR antibody (AA 150-268)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	ZFR
Binding Specificity:	AA 150-268
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZFR antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant Human Zinc finger RNA-binding protein (150-268AA)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	>95%, Protein G purified

Target Details

Target:	ZFR
Alternative Name:	ZFR (ZFR Products)
Background:	Background: Involved in postimplantation and gastrulation stages of development. Involved in the nucleocytoplasmic shuttling of STAU2. Binds to DNA and RNA (By similarity).

Target Details

Aliases: FLJ41312 antibody, hZFR antibody, M phase phosphoprotein homolog antibody, M-phase phosphoprotein homolog antibody, zfr antibody, ZFR_HUMAN antibody, zinc finger RNA binding protein antibody, Zinc finger RNA-binding protein antibody

UniProt: [Q96KR1](#)

Application Details

Application Notes: Recommended dilution: IF:1:50-1:200,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Immunofluorescence

Image 1. Immunofluorescence staining of HepG2 cells with ABIN7176651 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).