

Datasheet for ABIN7177115
anti-MMP13 antibody (AA 105-472) (FITC)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	MMP13
Binding Specificity:	AA 105-472
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MMP13 antibody is conjugated to FITC
Application:	Please inquire

Product Details

Immunogen:	Recombinant Mouse Collagenase 3 protein (105-472AA)
Isotype:	IgG
Cross-Reactivity:	Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	MMP13
Alternative Name:	Mmp13 (MMP13 Products)
Background:	Background: Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II

Target Details

and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CTGF. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CTGF. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion.

Aliases: Mmp13 antibody, Collagenase 3 antibody, EC 3.4.24.- antibody, Matrix metalloproteinase-13 antibody, MMP-13 antibody

UniProt: [P33435](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.