

Datasheet for ABIN7177415

anti-M-CSF/CSF1 antibody (AA 33-262) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	M-CSF/CSF1 (CSF1)
Binding Specificity:	AA 33-262
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This M-CSF/CSF1 antibody is conjugated to Biotin
Application:	ELISA

Product Details

Immunogen:	Recombinant Mouse Macrophage colony-stimulating factor 1 protein (33-262AA)
Isotype:	IgG
Cross-Reactivity:	Mouse
Purification:	>95%, Protein G purified

Target Details

Target:	M-CSF/CSF1 (CSF1)
Alternative Name:	Csf1 (CSF1 Products)
Background:	Background: Cytokine that plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes, such as

Target Details

macrophages and monocytes. Promotes the release of proinflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. Plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone development. Required for normal male and female fertility. Promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration. Plays a role in lipoprotein clearance.

Aliases: Csf1 antibody, Csf mMacrophage colony-stimulating factor 1 antibody, CSF-1 antibody, MCSF) [Cleaved into: Processed macrophage colony-stimulating factor 1] antibody

UniProt: [P07141](#)

Pathways: [RTK Signaling](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300
Constituents: 50 % Glycerol, 0.01M PBS, PH 7.4

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.