antibodies - online.com







anti-FYN antibody (Tyr530)

Images



Overview

Quantity:	100 μL
Target:	FYN
Binding Specificity:	Tyr530
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FYN antibody is un-conjugated
Application:	Western Blotting (WB), ELISA
Product Details	
Immunogen:	Synthesized non-phosphopeptide derived from Human Fyn around the phosphorylation site of tyrosine 530 (P-Q-Y(p)-Q-P).
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Target Details	
Target:	FYN
Alternative Name:	FYN (FYN Products)

Background:

Background: Non-receptor tyrosine-protein kinase that plays a role in many biological processes including regulation of cell growth and survival, cell adhesion, integrin-mediated signaling, cytoskeletal remodeling, cell motility, immune response and axon guidance. Inactive FYN is phosphorylated on its C-terminal tail within the catalytic domain. Following activation by PKA, the protein subsequently associates with PTK2/FAK1, allowing PTK2/FAK1 phosphorylation, activation and targeting to focal adhesions. Involved in the regulation of cell adhesion and motility through phosphorylation of CTNNB1 (beta-catenin) and CTNND1 (deltacatenin). Regulates cytoskeletal remodeling by phosphorylating several proteins including the actin regulator WAS and the microtubule-associated proteins MAP2 and MAPT. Promotes cell survival by phosphorylating AGAP2/PIKE-A and preventing its apoptotic cleavage. Participates in signal transduction pathways that regulate the integrity of the glomerular slit diaphragm (an essential part of the glomerular filter of the kidney) by phosphorylating several slit diaphragm components including NPHS1, KIRREL and TRPC6. Plays a role in neural processes by phosphorylating DPYSL2, a multifunctional adapter protein within the central nervous system, ARHGAP32, a regulator for Rho family GTPases implicated in various neural functions, and SNCA, a small pre-synaptic protein. Participates in the downstream signaling pathways that lead to T-cell differentiation and proliferation following T-cell receptor (TCR) stimulation. Also participates in negative feedback regulation of TCR signaling through phosphorylation of PAG1, thereby promoting interaction between PAG1 and CSK and recruitment of CSK to lipid rafts. CSK maintains LCK and FYN in an inactive form. Promotes CD28-induced phosphorylation of VAV1.

Kawakami T., Mol. Cell. Biol. 6:4195-4201(1986).

Semba K., Proc. Natl. Acad. Sci. U.S.A. 83:5459-5463(1986).

Rigley K., J. Immunol. 154:1136-1145(1995)

Aliases: C syn protooncogene antibody, Fyn antibody, FYN oncogene related to SRC FGR YES antibody, FYN_HUMAN antibody, OKT3 induced calcium influx regulator antibody, P59 FYN antibody, p59-Fyn antibody, Protein tyrosine kinase fyn antibody, Proto oncogene tyrosine protein kinase fyn antibody, Proto-oncogene c-Fyn antibody, Proto-oncogene Syn antibody, Protooncogene Syn antibody, Src like kinase antibody, Src yes related novel gene antibody, Src-like kinase antibody, Src/yes related novel antibody, SYN antibody, Tyrosine kinase p59fyn T antibody, Tyrosine kinase p59fyn(T) antibody, Tyrosine-protein kinase Fyn antibody

UniProt:

P06241

Pathways:

JAK-STAT Signaling, TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling

Pathway, Neurotrophin Signaling Pathway, Feeding Behaviour, CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2, Activated T Cell Proliferation, Thromboxane A2 Receptor Signaling

Application Details

Application Notes:	WB:1:500-1:3000,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl,
	0.02 % sodium azide and 50 % glycerol.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Western Blotting

Image 1. Western blot analysis of extracts from HeLa cells (Lane 2), A549 cells (Lane 3) and HepG2 cells (Lane 4), using Fyn (Ab-530) antiobdy. The lane on the left is treated with synthesized peptide.



Western Blotting

Image 2. Western blot analysis of extracts from HUVEC cells, MCF-7 cells and 293 cells, using Fyn (Ab-530) antibody.