

Datasheet for ABIN7180060
anti-p53 antibody (Ser378)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	p53 (TP53)
Binding Specificity:	Ser378
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This p53 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Synthesized non-phosphopeptide derived from Human p53 around the phosphorylation site of serine 378 (S-T-S(p)-R-H).
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Target Details

Target:	p53 (TP53)
Alternative Name:	TP53 (TP53 Products)

Target Details

Background: Background: Acts as a tumor suppressor in many tumor types, induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis, the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkl1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seem to have to effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. 7 0 2 5

Zakut-Houri R., EMBO J. 4:1251-1255(1985).

Lamb P., Mol. Cell. Biol. 6:1379-1385(1986).

Harlow E., Mol. Cell. Biol. 5:1601-1610(1985).

Aliases: Antigen NY-CO-13 antibody, BCC7 antibody, Cellular tumor antigen p53 antibody, FLJ92943 antibody, LFS1 antibody, Mutant tumor protein 53 antibody, p53 antibody, p53 tumor suppressor antibody, P53_HUMAN antibody, Phosphoprotein p53 antibody, Tp53 antibody, Transformation related protein 53 antibody, TRP53 antibody, tumor antigen p55 antibody, Tumor protein 53 antibody, Tumor protein p53 antibody, Tumor suppressor p53 antibody

UniProt: [P04637](#)

Pathways: [p53 Signaling](#), [MAPK Signaling](#), [PI3K-Akt Signaling](#), [Apoptosis](#), [AMPK Signaling](#), [Chromatin Binding](#), [ER-Nucleus Signaling](#), [Positive Regulation of Endopeptidase Activity](#), [Hepatitis C](#), [Protein targeting to Nucleus](#), [Autophagy](#), [Warburg Effect](#)

Application Details


Application Notes: WB:1:500-1:3000,

Restrictions: For Research Use only

Handling


Format:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Western Blotting

Image 1. Western blot analysis of extracts from HepG2 cells (Lane 2), using p53 (Ab-378) antibody. The lane on the left is treated with synthesized peptide.



Western Blotting

Image 2. Western blot analysis of extracts from 293 cells, using p53 (Ab-378) antibody.