antibodies - online.com







anti-CYP2D6 antibody (C-Term)





\sim							
	1//	\Box	$r \setminus$	/ [\bigcirc	1	٨,

Quantity:	100 μL
Target:	CYP2D6
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP2D6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Synthesized peptide derived from C-terminal of Human CYP2D6.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	

Target Details

Target:	CYP2D6
Alternative Name:	CYP2D6 (CYP2D6 Products)
Background: Background: Responsible for the metabolism of many drugs and environmental che	

it oxidizes. It is involved in the metabolism of drugs such as antiarrhythmics, adrenoceptor antagonists, and tricyclic antidepressants.

Gonzalez F.J., Genomics 2:174-179(1988).

Gonzalez F.J., Nature 331:442-446(1988).

Kimura S., Am. J. Hum. Genet. 45:889-904(1989).

Aliases: Cholesterol 25-hydroxylase antibody, CP2D6_HUMAN antibody, CPD6 antibody, CYP2D antibody, CYP2D6 antibody, CYP2D7AP antibody, CYP2D7BP antibody, CYP2D7P2 antibody, CYP2D8P2 antibody, CYP2DL1 antibody, CYPIID6 antibody, Cytochrome P450 2D6 antibody, Cytochrome P450 DB1 antibody, Cytochrome P450 family 2 subfamily D member 6 antibody, Cytochrome P450 family 2 subfamily D polypeptide 6 antibody, Cytochrome P450-DB1 antibody, Debrisoquine 4 hydroxylase antibody, Debrisoquine 4-hydroxylase antibody, Flavoprotein linked monooxygenase antibody, Microsomal monooxygenase antibody, P450 DB1 antibody, P450C2D antibody, P450DB1 antibody, Xenobiotic monooxygenase antibody

UniProt:

P10635

Application Details

Application Notes:	WB:1:500-1:3000,
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Format:

Liquid

Buffer:

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.

Preservative:

Sodium azide


This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage:

-20 °C,-80 °C

Storage Comment:

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western blot analysis of extracts from K562 cells, using CYP2D6 antibody.