# antibodies -online.com





Datasheet for ABIN718082

# anti-OSBP2 antibody (AA 221-320) (Cy7)



( )	11/0	K\ /	iew	1
	$\cup$	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	OSBP2
Binding Specificity:	AA 221-320
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OSBP2 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human OSBP2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	OSBP2
Alternative Name:	OSBP2 (OSBP2 Products)

## **Target Details**

Storage Comment:

Expiry Date:

rarget Details		
Background:	Synonyms: ORP 4, ORP4, OSBLP4, OSBP-related protein 4, OSBP2, OSBPL1, OSBPL4, Oxysterol	
	Binding Protein 2, Oxysterol Binding Protein-like 1, oxysterol binding protein-related protein 4,	
	OSBP2_HUMAN.	
	Background: The Oxysterol-binding protein (OSBP) family of proteins consist of OSBP (OSBP1)	
	and OSBP2 (ORP-4), which share a high overall similarity. OSBPs are involved in lipid	
	metabolism and signal transduction, as well as vesicle transport, and can translocate to the	
	periphery of Golgi membranes when they are bound to oxysterols. The OSBP protein transports	
	sterols from lysosomes to the nucleus, where sterols downregulate the genes for HMG	
	synthetase, HMG-CoA reductase and the low density lipoprotein receptor (LDLR). OSBP	
	localizes to the cytosol and is widely expressed, while OSBP2 is mainly detected in testis, retina	
	and fetal liver. The extracellular signal-regulated kinase (ERK) signaling pathway is controlled by	
	OSBP via its cholesterol-binding properties. OSBP binds with a high affinity to 25-hydroxy-	
	cholesterol (25-HC), a suppressor of cholesterol synthesis gene transcription in cultured cells.	
Gene ID:	23762	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw evoles	

12 months

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.