

Datasheet for ABIN7181491  
**anti-HIST1H2AG antibody (AA 86-98)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	HIST1H2AG
Binding Specificity:	AA 86-98
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HIST1H2AG antibody is un-conjugated
Application:	ELISA, Immunocytochemistry (ICC)

## Product Details

Immunogen:	Synthesized peptide derived from Human Histone H2A type 1 protein (86-98aa)
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Antigen Affinity Purified

## Target Details

Target:	HIST1H2AG
Alternative Name:	HIST1H2AG ( <a href="#">HIST1H2AG Products</a> )
Background:	Background: Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a

## Target Details

template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Aliases: Histone H2A type 1 (H2A.1) (Histone H2A/ptl), HIST1H2AG, HIST1H2AI, HIST1H2AK, HIST1H2AL, HIST1H2AM, H2AFP, H2AFC, H2AFD, H2AFI, H2AFN

UniProt: [P0C0S8](#)

## Application Details

Application Notes: Recommended dilution: ICC:1:10-1:100,

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Preservative: 0.03 % Proclin 300  
Constituents: 50 % Glycerol, 0.01M PBS, pH 7.4


Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

## Images



### Immunocytochemistry

**Image 1.** Immunocytochemistry analysis of ibHU diluted at 1:10 and staining in HeLa cells (treated with 30 mM sodium butyrate for 4h) performed on a Leica Bond™ system. The cells were fixed in 4 % formaldehyde, permeabilized using 0.2 % Triton X-100 and blocked with 10 % normal goat serum 30 min at RT. Then primary antibody (1 % BSA) was incubated at 4 °C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an

HRP conjugated SP system.