antibodies .- online.com





anti-TNFAIP8L2 antibody





Go to Product page

\sim							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

Quantity:	100 μL
Target:	TNFAIP8L2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNFAIP8L2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Synthesized peptide derived from internal of Human TNFAIP8L2.	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	

Target Details

Target:	TNFAIP8L2
Alternative Name:	TNFAIP8L2 (TNFAIP8L2 Products)
Background:	Background: Acts as a negative regulator of innate and adaptive immunity by maintaining
	immune homeostasis. Negative regulator of Toll-like receptor and T-cell receptor function.
	Prevents hyperresponsiveness of the immune system and maintains immune homeostasis.

Inhibits JUN/AP1 and NF-kappa-B activation. Promotes Fas-induced apoptosis By similarity.

Chen Y.H., Cell 133:415-426(2008).

Sugano S., Nat. Genet. 36:40-45(2004).

Schupp I., BMC Genomics 8:399-399(2007)

Aliases: AW610835 antibody, FLJ23467 antibody, Inflammation factor 20 antibody, Inflammation factor protein 20 antibody, TIPE2 antibody, TNF alpha-induced protein 8-like protein 2 antibody, TNFAIP8-like protein 2 antibody, TNFAIP8L2 antibody, TP8L2_HUMAN antibody, Tumor necrosis factor alpha induced protein 8 like 2 antibody, Tumor necrosis factor alpha-induced protein 8-like protein 2 antibody

UniProt:


Q6P589

Application Details

Application Notes:	WB:1:500-1:3000,
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02 % sodium azide and 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western blot analysis of extracts from HuvEc cells, using TNFAIP8L2 antibody.