



[Go to Product page](#)

Datasheet for ABIN718487  
**anti-APEX2 antibody (AA 301-400) (Cy7)**

### Overview

Quantity:	100 µL
Target:	APEX2
Binding Specificity:	AA 301-400
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This APEX2 antibody is conjugated to Cy7
Application:	Western Blotting (WB)

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human APEX2
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

### Target Details

Target:	APEX2
Alternative Name:	APEX2 ( <a href="#">APEX2 Products</a> )
Background:	Synonyms: APE2, XTH2, ZGRF2, APEXL2, DNA-(apurinic or apyrimidinic site) lyase 2, AP

## Target Details

---

endonuclease XTH2, APEX nuclease 2, APEX nuclease-like 2, Apurinic-apyrimidinic endonuclease 2, AP endonuclease 2, APEX2

Background: Function as a weak apurinic/apyrimidinic (AP) endodeoxyribonuclease in the DNA base excision repair (BER) pathway of DNA lesions induced by oxidative and alkylating agents. Initiates repair of AP sites in DNA by catalyzing hydrolytic incision of the phosphodiester backbone immediately adjacent to the damage, generating a single-strand break with 5'-deoxyribose phosphate and 3'-hydroxyl ends. Displays also double-stranded DNA 3'-5' exonuclease, 3'-phosphodiesterase activities. Shows robust 3'-5' exonuclease activity on 3'-recessed heteroduplex DNA and is able to remove mismatched nucleotides preferentially. Shows fairly strong 3'-phosphodiesterase activity involved in the removal of 3'-damaged termini formed in DNA by oxidative agents. In the nucleus functions in the PCNA-dependent BER pathway. Required for somatic hypermutation (SHM) and DNA cleavage step of class switch recombination (CSR) of immunoglobulin genes. Required for proper cell cycle progression during proliferation of peripheral lymphocytes.

---

Gene ID: 27301

UniProt: [Q9UBZ4](#)

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Storage: -20 °C

---

## Handling

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months