antibodies -online.com







anti-LIF antibody (C-Term)





Overview

Quantity:	100 μg
Target:	LIF
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LIF antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Synthesized peptide derived from the C-terminal region of Human LIF.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Target Details

Target:	LIF
Alternative Name:	LIF (LIF Products)
Background:	CDF antibody, Cholinergic Differentiation Factor antibody, D factor antibody, DIA antibody,

Target Details

Differentiation inducing factor antibody, differentiation inhibitory activity antibody,
Differentiation stimulating factor antibody, Differentiation-stimulating factor antibody,
Emfilermin antibody, Hepatocyte stimulating factor III antibody, HILDA antibody, Human
interleukin in DA cells antibody, Leukemia inhibitory factor antibody, LIF antibody, LIF_HUMAN
antibody, Melanoma derived LPL inhibitor antibody, Melanoma-derived LPL inhibitor antibody,
MLPLI antibody

UniProt:

P15018

Pathways:


JAK-STAT Signaling, Positive Regulation of Peptide Hormone Secretion, Negative Regulation of Hormone Secretion, Stem Cell Maintenance, Growth Factor Binding

Application Details

Application Notes:	WB:1:500-1:2000, ELISA:1:20000,
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western Blot analysis of K562 cells using LIF Polyclonal Antibody