# antibodies - online.com







## anti-MAG antibody (C-Term)



Image



0	1 /	-	K	/1	-	1 A
u	\/	$\vdash$	I \	/ I	$\vdash$	1/1

Quantity:	100 μg
Target:	MAG
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAG antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

#### **Product Details**

Immunogen:	Synthesized peptide derived from the C-terminal region of Human MAG.	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.	

#### **Target Details**

Target:	MAG
Alternative Name:	MAG (MAG Products)
Background:	GMA antibody, MAG antibody, MAG_HUMAN antibody, Myelin associated glycoprotein antibody,

#### **Target Details**

Myelin-associated glycoprotein antibody, S MAG antibody, S-MAG antibody, Sialic acid bine	ding
lg like lectin 4A antibody, Sialic acid binding immunoglobulin like lectin 4A antibody, Siglec	4a
antibody, Siglec-4a antibody, SIGLEC4A antibody, SPG75 antibody	

UniProt: P20916

Pathways: Neurotrophin Signaling Pathway

#### **Application Details**


Application Notes:	WB:1:500-1:2000, ELISA:1:20000,

Restrictions: For Research Use only

### Handling

Format:	Liquid
Buffer:	Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

#### **Images**



#### **Western Blotting**

**Image 1.** Western Blot analysis of HeLa HepG2 cells using MAG Polyclonal Antibody