

Datasheet for ABIN7185512
anti-SDHA antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	SDHA
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SDHA antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Synthesized peptide derived from the C-terminal region of Human SDHA.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Target Details

Target:	SDHA
Alternative Name:	SDHA (SDHA Products)
Background:	CMD1GG antibody, DHSA_HUMAN antibody, Flavoprotein subunit of complex II antibody, Fp

Target Details

antibody, PGL5 antibody, SDH 2 antibody, SDH1 antibody, SDH2 antibody, SDHA antibody, SDHF antibody, Succinate dehydrogenase [ubiquinone] flavoprotein subunit antibody, Succinate dehydrogenase [ubiquinone] flavoprotein subunit mitochondrial antibody, Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial antibody, Succinate dehydrogenase complex flavoprotein subunit A antibody, Succinate dehydrogenase complex flavoprotein subunit antibody, Succinate dehydrogenase complex flavoprotein subunit precursor antibody, Succinate dehydrogenase complex subunit A antibody, Succinate dehydrogenase complex subunit A flavoprotein (Fp) antibody, Succinate dehydrogenase complex subunit A flavoprotein antibody

UniProt: [P31040](#)

Application Details

Application Notes: WB:1:500-1:2000, ELISA:1:10000,

Restrictions: For Research Use only

Handling

Format: Liquid


Buffer: Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: -20 °C,-80 °C

Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western Blot analysis of NIH-3T3 Jurkat K562 HepG2 RAT-KIDNEY cells using SDHA Polyclonal Antibody



Western Blotting

Image 2. Western Blot analysis of HepG2 cells using SDHA Polyclonal Antibody