

Datasheet for ABIN718565
anti-HSD17B6 antibody (AA 61-160) (HRP)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HSD17B6
Binding Specificity:	AA 61-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSD17B6 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HSD17B6
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	HSD17B6
Alternative Name:	HSD17B6 (HSD17B6 Products)

Target Details

Background:	Synonyms: 17-beta-HSD 6, 17-beta-HSD6, 17-beta-hydroxysteroid dehydrogenase type 6, 3 hydroxysteroid epimerase, 3alpha >beta hydroxysteroid epimerase, 3alpha >beta hydroxysteroid epimerasel, 3-alpha->beta-HSE, 3-alpha->beta-hydroxysteroid epimerase, H17B6_HUMAN, HSD17B6, HSE, Hydroxysteroid 17 beta dehydrogenase 6, Hydroxysteroid 17 beta dehydrogenase 6 homolog mouse, Hydroxysteroid 17 beta dehydrogenase 6, NAD+ dependent 3 alpha hydroxysteroid dehydrogenase 3 hydroxysteroid epimerase, NAD+ dependent 3 alpha hydroxysteroid dehydrogenase, Oxidative 3 alpha hydroxysteroid dehydrogenase, Oxidative 3-alpha hydroxysteroid dehydrogenase, Oxidoreductase, Retinol dehydrogenase, RODH, SDR9C6,Short chain dehydrogenase/reductase family 9C, member 6. Background: NAD-dependent oxidoreductase with broad substrate specificity that shows both oxidative and reductive activity (in vitro). Has 17-beta-hydroxysteroid dehydrogenase activity towards various steroids (in vitro). Converts 5-alpha-androstan-3-alpha,17-beta-diol to androsterone and estradiol to estrone (in vitro). Has 3-alpha-hydroxysteroid dehydrogenase activity towards androsterone (in vitro). Has retinol dehydrogenase activity towards all-trans-retinol (in vitro). Can convert androsterone to epi-androsterone. Androsterone is first oxidized to 5-alpha-androstane-3,17-dione and then reduced to epi-andosterone. Can act on both C-19 and C-21 3-alpha-hydroxysteroids.Tissue specificity, Detected in liver and prostate (at protein level). Detected in adult liver, lung, brain, placenta, prostate, adrenal gland, testis, mammary gland, spleen, spinal cord and uterus. Detected in caudate nucleus, and at lower levels in amygdala, corpus callosum, hippocampus, substantia nigra and thalamus. Detected in fetal lung, liver and brain.Sequence similarities, Belongs to the short-chain dehydrogenases/reductases (SDR) family.
Gene ID:	8630
Pathways:	Steroid Hormone Biosynthesis

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
---------	--------

Handling

Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months