
antibodies -online.com





anti-WTAP antibody (C-Term)





Go to Product page

\sim							
	1//	$rac{1}{2}$	`\/	L	Р	1	٨

Quantity:	100 μg
Target:	WTAP
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WTAP antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Synthesized peptide derived from the C-terminal region of Human WTAP.
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Target Details

Target:	WTAP
Alternative Name:	WTAP (WTAP Products)
Background:	DKFZp686F20131 antibody, Female-lethal(2)D homolog antibody, FL2D_HUMAN antibody,

hFL(2)D antibody, KIAA0105 antibody, MGC3925 antibody, Mum2 antibody, OTTHUMP00000017522 antibody, OTTHUMP00000017523 antibody, PNAS 132 antibody, PNAS-132 antibody, PNAS-132 antibody, Pre-mRNA-splicing regulator WTAP antibody, Pre-mRNA-splicing regulator WTAP antibody, Putative pre mRNA splicing regulator female lethal 2D homolog antibody, Putative pre mRNA splicing regulator female lethal(2D) antibody, Putative pre-mRNA splicing regulator female lethal(2D) homolog antibody, putative pre-mRNA splicing regulator female-lethal(2D) antibody, Wilms tumor 1 associated protein antibody, Wilms tumor 1 associating protein antibody, Wilms' tumor 1 associating protein antibody, Wilms' tumour 1-associating protein antibody, WT1 associated protein antibody, WT1-associated protein antibody, wtap antibody

UniProt:


Q15007

Application Details

Application Notes:	WB:1:500-1:2000, IHC:1:100-1:300, ELISA:1:20000,
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Western Blotting

Image 1. Western Blot analysis of HuvEc cells using WTAP Polyclonal Antibody