# antibodies .- online.com





# anti-SH2D2A antibody (Internal Region)



Image



Go to Product page

### Overview

Quantity:	100 μg
Target:	SH2D2A
Binding Specificity:	Internal Region
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH2D2A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

### **Product Details**

Immunogen:	Synthesized peptide derived from the Internal region of Human Lad.
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

# **Target Details**

Target:	SH2D2A
Alternative Name:	SH2D2A (SH2D2A Products)
Background:	F2771 antibody, SCAP antibody, SH2 domain containing adapter protein antibody, SH2 domain

#### **Target Details**

containing protein 2A antibody, SH2 domain protein 2A antibody, SH2 domain-containing adapter protein antibody, SH2 domain-containing protein 2A antibody, SH22A\_HUMAN antibody, Sh2d2a antibody, T cell specific adapter protein antibody, T cell specific adapter protein TSAd antibody, T cell-specific adapter protein antibody, T cell-specific adapter protein antibody, T lymphocyte specific adaptor protein antibody, TSAd antibody, VEGF receptor associated protein antibody, VEGF receptor-associated protein antibody, VRAP antibody

UniProt: Q9NP31

Pathways: Signaling Events mediated by VEGFR1 and VEGFR2, VEGF Signaling


# **Application Details**

Application Notes: WB:1:500-1:2000, IHC:1:100-1:300, ELISA:1:10000,

Restrictions: For Research Use only

### Handling

Format:	Liquid
Buffer:	Liquid in PBS containing 50 % glycerol, 0.5 % BSA and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



## **Western Blotting**

**Image 1.** Western Blot analysis of HepG2 cells using Lad Polyclonal Antibody