antibodies - online.com







anti-CCL23 antibody (C-Term)





\sim	
$() \setminus (\cap)$	$r_1/1011/1$
\cup	rview
	_

Quantity:	100 μL
Target:	CCL23
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Synthetic peptide corresponding to residues near the C terminal of Human C-C motif chemokine 6
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Antigen Affinity Purified

Target Details

Target:	CCL23
Alternative Name:	CCL23 (CCL23 Products)
Background:	Background: In mice, CCL6 is expressed in cells from neutrophil and macrophage lineages, and can be greatly induced under conditions suitable for myeloid cell differentiation. It is highly expressed in bone marrow cultures that have been stimulated with the cytokine GM-CSF. Some

low levels of gene expression also occur in certain cell lines of myeloid origin (e.g. the immature myeloid cell lines DA3 and 32D cl3, and the macrophage cell line P388D) that can also be greatly induced in culture with GM-CSF. However, in activated T cell lines, expression of CCL6 is greatly reduced. CCL6 can also be induced in the mouse lung by the cytokine interleukin 13 Aliases: C C motif chemokine ligand 23 antibody, C-C motif chemokine 23 antibody, C6 beta chemokine antibody, CCL 23 antibody, CCL23 antibody, CCL23(30-99) antibody, CCL23_HUMAN antibody, Chemokine (C C motif) ligand 23 antibody, Chemokine CC motif ligand 23 antibody, ck beta 8 antibody, CK-beta-8 antibody, CKB 8 antibody, CKB-8 antibody, CKb-8-1 antibody, CKB8 antibody, Hmrp-2a antibody, Macrophage inflammatory protein 3 antibody, MIP 3 antibody, MIP-3 antibody, MIP3 antibody, MPIF 1 antibody, MPIF-1 antibody, MPIF-1 antibody, MPIF-1 antibody, SCYA 23 antibody, Small inducible cytokine A 23 antibody, Small inducible cytokine subfamily A, member 23 antibody, Small-inducible cytokine A 23 antibody

UniProt:


P55773

Application Details

Application Notes:	IHC:1:50-1:100,
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Rabbit IgG in pH 7.3 PBS, 0.05 % Sodium azide, 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.



Immunohistochemistry

Image 1. Immunohistochemical analysis of paraffinembedded Human lung cancer tissue using at dilution 1/50.