

Datasheet for ABIN7189479
anti-HMMR antibody (AA 280-340)[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	HMMR
Binding Specificity:	AA 280-340
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMMR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA

Product Details

Immunogen:	Synthetic peptide from Human protein at AA range: 280-340
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.

Target Details

Target:	HMMR
Alternative Name:	HMMR (HMMR Products)
Background:	CD168 antibody, CD168 antigen antibody, HMMR antibody, HMMR_HUMAN antibody,

Target Details

Hyaluronan mediated motility receptor antibody, Hyaluronan-mediated motility receptor (RHAMM) antibody, IHABP antibody, Intracellular hyaluronic acid-binding protein antibody, MGC119494 antibody, MGC119495 antibody, OTTHUMP00000196920 antibody, Receptor for hyaluronan-mediated motility antibody, RHAMM antibody

UniProt: [O75330](#)

Pathways: [Glycosaminoglycan Metabolic Process](#)

Application Details

Application Notes: WB:1:500-2000, IHC:1:50-300, ELISA:1:10000-20000,

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: PBS, pH 7.4, containing 0.02 % sodium azide as Preservative and 50 % Glycerol.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C,-80 °C


Storage Comment: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images




Immunohistochemistry

Image 1. Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200



Western Blotting

Image 2. Western blot analysis of 293T lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000



Immunohistochemistry

Image 3. Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200