antibodies.com

Datasheet for ABIN7192738 anti-TBC1D21 antibody

Image



Overview

Quantity:	100 µL
Target:	TBC1D21
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBC1D21 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Synthetic peptide of Human TBC1D21
Isotype:	lgG
Cross-Reactivity:	Human
Purification:	Antigen affinity purification


Target Details

Target:	TBC1D21
Alternative Name:	TBC1D21 (TBC1D21 Products)
Background:	Background: TBC1D21 gene (TBC1 domain family member 21) is located on human chromosome 15, acts as a GTPase-activating protein for Rab family proteins. Rab family
	proteins comprise G proteins of the Ras superfamily that belong to a protein superfamily of small GTPases. Approximately 70 types of Rabs have been identified in humans. They are

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7192738 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
	regulators of membrane traffic, including vesicle formation, vesicle movement along actin and tubulin networks, and membrane fusion. Aliases: TBC1D21 antibody, TBC1 domain family member 21 antibody
UniProt:	Q8IYX1
Application Details	
Application Notes:	ELISA:1:1000-1:2000, WB:1:200-1:1000,
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	-20 °C, pH 7.4 PBS, 0.05 % Sodium azide, 40 % Glycerol
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,-80 °C
Storage Comment:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Images



Western Blotting

Image 1. Gel: 8 % SDS-PAGE, Lysate: 40 µg, Lane: Hepg2 cells, Primary antibody: ABIN7192738(TBC1D21 Antibody) at dilution 1/300, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds