

Datasheet for ABIN719398

anti-ADORA1 antibody (AA 225-326) (Biotin)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	ADORA1	
Binding Specificity:	AA 225-326	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ADORA1 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Adenosine A1 Receptor	
	IgG	
Isotype:	igo	
Cross-Reactivity:	Human, Mouse	
Cross-Reactivity:	Human, Mouse	
Cross-Reactivity: Predicted Reactivity:	Human, Mouse Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit	
Cross-Reactivity: Predicted Reactivity: Purification:	Human, Mouse Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit	
Cross-Reactivity: Predicted Reactivity: Purification: Target Details	Human, Mouse Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit Purified by Protein A.	
Cross-Reactivity: Predicted Reactivity: Purification: Target Details Target:	Human, Mouse Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit Purified by Protein A. ADORA1	

Background: Adenosine is involved in a variety of processes, including the synthesis of urea, the anti-inflammatory response, and the inhibition of protein synthesis. The Adenosine receptors, including Adenosine A1-R, Adenosine A2A-R, Adenosine A2B-R and Adenosine A3-R, are integral membrane proteins that are members of the G protein-coupled receptor family. Adenosine A1-R mediates ureagenesis in a partially calcium-dependent manner. Adenosine is known to mediate coronary vasodilation via Adenosine A2A-R. Collagen synthesis and total protein synthesis are inhibited in certain cells by Adenosine, acting via the A2B receptors. Activation of Adenosine A3-R inhibits the induction of TNF?and blocks the endotoxin CD14 receptor signal transduction pathway.

Gene ID: 134

UniProt: P30542

Pathways: EGFR Signaling Pathway, Negative Regulation of Hormone Secretion, Synaptic Membrane

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months