antibodies -online.com







ACVRL1 Protein (His tag,Fc Tag)





Overview


Quantity:	200 μg
Target:	ACVRL1
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This ACVRL1 protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Rat ALK-1/ACVRL1 Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Ala 118
Characteristics:	A synthetic DNA sequence encoding the rat ACVRL1 (NP_071886.1) extracellular domain (Met 1-Ala 118) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	$<$ 1.0 EU per μg of the protein as determined by the LAL method
Biological Activity Comment:	Measured by its ability to inhibit BMP9-induced alkaline phosphatase production by MC3T3E1 mouse chondrogenic cells. David, L. et al. (2007) Blood 109:1953. The ED50 for this effect is typically 40-200 ng/mL in the presence of 2 ng/mL of recombiant human BMP9.

Target Details

Target:	ACVRL1
Alternative Name:	ALK-1/ACVRL1 (ACVRL1 Products)
Background:	Background: Activin A receptor, type II-like 1 (ACVRL1), also known as ALK-1 (activin receptor
	like kinase 1), is an endothelial-specific type I receptor of the TGF-beta (transforming growth
	factor beta) receptor family of ligands. On ligand binding, a heteromeric receptor complex
	forms consisting of two type II and two type I transmembrane serine/threonine kinases.
	ACVRL1 protein is expressed in certain blood vessels of kidney, spleen, heart and intestine,
	serving as an important role during vascular development. Mutations in ACVRL1 gene are
	associated with hemorrhagic telangiectasia type 2, also known as Rendu-Osler-Weber
	syndrome 2 and vascular disease.
	Synonym: R3,R-3,SETHKIR,MGC91691,Acvrl1
Molecular Weight:	38.7 kDa
NCBI Accession:	NP_071886
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.