

Datasheet for ABIN7194682

**Cathepsin L Protein (His tag)**[Go to Product page](#)

## Overview

|                               |  |
|-------------------------------|--|
| Quantity:                     | 50 µg  |
| Target:                       | Cathepsin L (CTSL1)                                |
| Origin:                       | Human  |
| Source:                       | HEK-293 Cells                                      |
| Protein Type:                 | Recombinant  |
| Biological Activity:          | Active   |
| Purification tag / Conjugate: | This Cathepsin L protein is labelled with His tag. |

## Product Details

|                              |  |
|------------------------------|--|
| Purpose:                     | Recombinant Human Cathepsin L/CTSL Protein (His Tag)(Active)   |
| Sequence:                    | Met 1-Val 333  |
| Characteristics:             | A DNA sequence encoding the pro form of human Cathepsin-L1 (NP_001903.1) (Met 1-Val 333) was expressed, fused with a polyhistidine tag at the C-terminus.                      |
| Purity:                      | > 90 % as determined by reducing SDS-PAGE.   |
| Endotoxin Level:             | < 1.0 EU per µg as determined by the LAL method.   |
| Biological Activity Comment: | Measured by its binding ability in a functional ELISA. Immobilized human CD74 at 5 µg/ml (100 µl/well) can bind biotinylated human CTSL1 with a linear range of 3.2-400 ng/ml. |

## Target Details

|         |                     |
|---------|---------------------|
| Target: | Cathepsin L (CTSL1) |
|---------|---------------------|

## Target Details

|                   |   |
|-------------------|---|
| Alternative Name: | Cathepsin L/CTSL ( <a href="#">CTSL1 Products</a> )   |
| Background:       | <p>Background: Cathepsin L is a lysosomal cysteine protease that plays a major role in intracellular protein catabolism, and is potent in degrading collagen, laminin, elastin, as well as alpha-1 protease inhibitor and other structural proteins of basement membranes. It is secreted by liver flukes at all stages of their development in the mammalian host, are believed to play important roles in facilitating parasite migration (tissue degradation), feeding and immuno-evasion. Like many proteases, Cathepsin L is synthesized as an inactive preproenzyme, and cleavage of the 96-residue proregion is necessary to generate the fully active 221-residue mature enzyme. Studies have demonstrated that cleavage of the proregion occur autocatalytically under acidic conditions. The enzyme takes part in nutrient acquisition by catabolizing host proteins to absorbable peptides, facilitates the migration of the parasite through the host intestine and liver by cleaving interstitial matrix proteins such as fibronectin, laminin and native collagen and is implicated in the inactivation of host immune defenses by cleaving immunoglobulins. Recently, Cathepsin L has been shown to suppress Th1 immune response in infected laboratory animals making them susceptible to concurrent bacterial infections. Cathepsin L is synthesized in large amounts and secreted by many malignantly transformed cells, and induced by growth factors and tumor promoters. In addition to its role in protein degradation, evidence has accumulated for the participation of Cathepsin L in various physiological and pathological processes, such as tumor invasion and metastasis, bone resorption, spermatogenesis, and arthritis. Accordingly, Cathepsin L may prove useful as a diagnostic or prognostic marker of human tumor malignancy.</p> <p>Synonym: Cathepsin L1, Major Excreted Protein, MEP, CTSL1, CTSL</p> |
| Molecular Weight: | 37.3 kDa  |
| NCBI Accession:   | <a href="#">NP_001903</a>   |
| Pathways:         | <a href="#">Activation of Innate immune Response</a> , <a href="#">Toll-Like Receptors Cascades</a>   |

## Application Details

|                 |  |
|-----------------|--|
| Restrictions:   | For Research Use only  |
| Handling        |  |
| Format:         | Lyophilized  |
| Reconstitution: | Please refer to the printed manual for detailed information. |
| Buffer:         | Lyophilized from sterile 50 mM NaAc, 0.1M NaCl, pH 5.0       |

## Handling

---

Storage: 4 °C, -20 °C, -80 °C

---

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.