

Datasheet for ABIN719473 anti-ADA antibody (AA 65-170) (Biotin)



Overview

Alternative Name:

Quantity:	100 μL
Target:	ADA
Binding Specificity:	AA 65-170
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADA antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Adenosine deaminase
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Pig,Horse
	Human, Rat, Cow, Fig. Horse
Purification:	Purified by Protein A.
Purification: Target Details	

Adenosine deaminase (ADA Products)

Target Details

Background:	Synonyms: Adenosine deaminase, Adenosine aminohydrolase, ADA, ADA1
	Background: Catalyzes the hydrolytic deamination of adenosine and 2-deoxyadenosine. Plays
	an important role in purine metabolism and in adenosine homeostasis. Modulates signaling by
	extracellular adenosine, and so contributes indirectly to cellular signaling events. Acts as a
	positive regulator of T-cell coactivation, by binding DPP4. Its interaction with DPP4 regulates
	lymphocyte-epithelial cell adhesion.
Gene ID:	100
UniProt:	P00813
Pathways:	Regulation of G-Protein Coupled Receptor Protein Signaling, Ribonucleoside Biosynthetic
	Process
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months