

Datasheet for ABIN7194949

CD80 Protein (CD80) (His tag)





Overview

Quantity:	100 μg
Target:	CD80
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD80 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse CD80/B7-1 Protein (His Tag)(Active)
Sequence:	Met 1-Lys 245
Characteristics:	A DNA sequence encoding the mouse CD80 (Q00609-1) extracellular domain (Met 1-Lys 245) was expressed, with a polyhistidine tag at the C-terminus.
Purity:	> 97 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to bind recombinant mouse CD28 in a functional ELISA.

Target Details

Target:	CD80
Alternative Name:	CD80/B7-1 (CD80 Products)

Target Details

Background:

Background: The B-lymphocyte activation antigen B7-1 (referred to as B7), also known as CD80, is a member of cell surface immunoglobulin superfamily and is expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. As costimulatory ligands, B7-1 which exists predominantly as dimer and the related protein B7-2, interact with the costimulatory receptors CD28 and cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) expressed on T cells, and thus constitute one of the dominant pathways that regulate T cell activation and tolerance, cytokine production, and the generation of CTL. The B7/CD28/CTLA4 pathway has the ability to both positively and negatively regulate immune responses. CD80 is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas.Immune CheckpointImmune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: IHC Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: WB AntibodiesImmune Checkpoint ProteinsImmune Checkpoint Targets Coinhibitory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy Targeted Therapy Synonym: T-lymphocyte activation antigen CD80; Activation B7-1 antigen; B7; CD80; Cd28l; Ly-53;Ly53;MIC17;TSA1

Molecular Weight:

25.1 kDa

Pathways:

TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Positive Regulation of Immune Effector Process, Cancer Immune Checkpoints


Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.