

Datasheet for ABIN7195018 Cell Adhesion Molecule 4 Protein (CADM4) (Fc Tag)



Overview	
Quantity:	50 µg
Target:	Cell Adhesion Molecule 4 (CADM4)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Cell Adhesion Molecule 4 protein is labelled with Fc Tag.
Product Details	
Purpose:	Recombinant Human CADM4/IGSF4C/NECL-4 Protein (Fc Tag)
Sequence:	Met 1-Tyr323
Characteristics:	A DNA sequence encoding the human CADM4 (NP_660339.1) (Met1-Tyr323) was expressed with the Fc region of human IgG1 at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per μ g as determined by the LAL method.
Target Details	
Target:	Cell Adhesion Molecule 4 (CADM4)
Alternative Name:	CADM4/IGSF4C/NECL-4 (CADM4 Products)
Background:	Background: Immunoglobulin superfamily member 4C (IGSF4C), also known as CADM4 or

NECL-4, is an immunoglobulin (Ig) superfamily molecule showing significant homology with a

lung tumor suppressor, TSLC1. CADM4/IGSF4C/NECL-4 protein is mainly expressed in the

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7195018 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

kidney, bladder, and prostate in addition to the brain. Experiments have reported the biological		
significance of CADM4/IGSF4C/NECL-4 in the urinary tissues. An immunohistochemical study		
reveals that CADM4 is expressed at the cell-cell attachment sites in the renal tubules, the		
transitional epithelia of the bladder, and the glandular epithelia of the prostate. IGSF4-		
immunoreactivity (IR) was observed diffusely in the telencephalic wall, whereas it became		
rather confined to the subplate, the cortical plate and the subventricular zone as the		
development proceeded. IGSF4-IR gradually decreased after birth and disappeared in		
adulthood. IGSF4 remained at low levels throughout embryonic stage, whereas it increased		
after birth. These spatiotemporal patterns of the expression suggest that IGSF4 plays crucial		
roles in the development of both telencephalon and cerebellum. CADM4/IGSF4C/NECL-4 is		
ectopically expressed in adult T-cell leukemia (ATL) cells, providing not only a diagnostic marker		
for ATL, but also a possible therapeutic target against its invasion. The distinct roles of		
CADM4/IGSF4C/NECL-4 in the oncogenesis of carcinomas and ATL could be due to tissue-		
specific differences in the downstream cascades, and is a novel concept with respect to cell		
adhesion in human oncogenesis.		

Synonym: IGSF4C,Necl-4,NECL4,synCAM4,TSLL2

Molecular Weight:	60.1 kDa
NCBI Accession:	NP_660339

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN7195018 | 07/25/2024 | Copyright antibodies-online. All rights reserved.