

## Datasheet for ABIN7195054 CCL22 Protein (His tag)



[Go to Product page](#)

### Overview

Quantity:	50 µg
Target:	CCL22
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This CCL22 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human CCL22/MDC Protein (His Tag)
Sequence:	Gly 25-Gln 93
Characteristics:	A DNA sequence encoding the secreted form of human CCL22 (000626) (Gly 25-Gln 93) was expressed, with a polyhistidine tag at the N-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.

### Target Details

Target:	CCL22
Alternative Name:	CCL22/MDC ( <a href="#">CCL22 Products</a> )
Background:	Background: Chemokine (C-C motif) ligand 22(ABCD-1 / CCL22) is a kind of CC chemokine which is a family of secreted proteins involved in immunoregulatory and inflammatory processes. The cytokine displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for

## Target Details

primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes. This ABCD-1 / CCL22 chemokine binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated / effector T lymphocytes to inflammatory sites and other aspects of activated T-lymphocyte physiology. ABCD-1 / CCL22 is highly expressed in macrophage and in monocyte-derived dendritic cells, and thymus, and in Langerhans' cell histiocytosis and atopic dermatitis but not in dermatopathic lymphadenopathy. This chemokine is also found in lymph node, appendix, activated monocytes, resting and activated macrophages. This protein is lower expressed in lung and spleen and very weakly expressed in small intestine.

Synonym: A-152E5.1,ABCD-1,DC/B-CK,MDC,SCYA22,STCP-1

Molecular Weight:	9.7 kDa
UniProt:	<a href="#">O00626</a>

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 30 % Acetonitrile, 0.1 % TFA
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.