



[Go to Product page](#)

Datasheet for ABIN7195545

EPH Receptor B1 Protein (EPHB1) (His tag)

1 Image

Overview

Quantity:	200 µg
Target:	EPH Receptor B1 (EPHB1)
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This EPH Receptor B1 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Mouse EphB1/EPHT2 Protein (His Tag)(Active)
Sequence:	Met 1-Leu 539
Characteristics:	A DNA sequence encoding the mouse EPHB1 (Q8CBF3-1) extracellular domain (Met 1-Leu 539) was expressed, with a polyhistidine tag at the C-terminus.
Purity:	> 98 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Immobilized mouse EPHB1-His at 10 µg/ml (100 µl/well) can bind mouse EFNB1-Fc, The EC50 of mouse EFNB1-Fc is 0.04-0.08 µg/ml.

Target Details

Target:	EPH Receptor B1 (EPHB1)
---------	-------------------------

Target Details

Alternative Name: EphB1/EPHT2 ([EPHB1 Products](#))

Background: Ephrin type-B receptor 1, also known as EphB1, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. EphB2 receptor tyrosine kinase phosphorylates syndecan-2 and that this phosphorylation event is crucial for syndecan-2 clustering and spine formation. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity⁶. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). Ligand-mediated activation of Ephs induce various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer. EphB receptor tyrosine kinases are enriched at synapses, suggesting that these receptors play a role in synapse formation or function. We find that EphrinB binding to EphB induces a direct interaction of EphB with NMDA-type glutamate receptors. This interaction occurs at the cell surface and is mediated by the extracellular regions of the two receptors, but does not require the kinase activity of EphB.

Synonym:

9330129L11,AW488255,C130099E04Rik,Cek6,Elk,Elkh,ENSMUSG00000074119,Hek6,Net

Molecular Weight: 59.6 kDa

Pathways: [RTK Signaling](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized


Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.