antibodies -online.com





EPH Receptor B6 Protein (EPHB6) (Fc Tag)



Overview

Quantity:	100 μg
Target:	EPH Receptor B6 (EPHB6)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This EPH Receptor B6 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Human EphB6 Protein (Fc Tag)(Active)
Sequence:	Met 1-Ser 579
Characteristics:	The extracellular domain (Met 1-Ser 579) of human EphB6 (NP_004436.1) precursor was fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	1. Measured by its binding ability in a functional ELISA.2. Immobilized recombinant human EphrinB1 at 10 μ g/ml (100 μ l/well) can bind human EphB6 with a linear range of 0.16-4 μ g/ml.3. Immobilized recombinant human EphrinB2 at 10 μ g/ml (100 μ l/well) can bind human EphB6 with a linear range of 1.28-32 ng/ml.

Target Details

Target:	EPH Receptor B6 (EPHB6)
Alternative Name:	EphB6 (EPHB6 Products)
Background:	Background: Ephrins are divided into the ephrin-A (EFNA) class and the ephrin-B (EFNB) class
	based on their structures and sequence relationships. Ephrin receptors make up the largest
	subgroup of the receptor tyrosine kinase (RTK) family. EphB6 is an unusual Eph receptor,
	lacking catalytic capacity due to alterations in its kinase domain. Interestingly, increased
	metastatic activity is associated with reduced EphB6 receptor expression in several tumor
	types, including breast cancer. This emphasizes the potential of EphB6 to act as a suppressor
	of cancer aggressiveness. EphB6 suppress cancer invasiveness through c-Cbl-dependent
	signaling, morphologic changes, and cell attachment and indicate that EphB6 may represent a
	useful prognostic marker and a promising target for therapeutic approaches. EphB6 can both
	positively and negatively regulate cell adhesion and migration, and suggest that tyrosine
	phosphorylation of the receptor by an Src family kinase acts as the molecular switch for the
	functional transition. In addition, Ephrin-B2 may be a physiological ligand for the EphB6
	receptor.
	Synonym: HEP
Molecular Weight:	86.5 kDa
NCBI Accession:	NP_004436
Pathways:	RTK Signaling, Hormone Transport
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.