

[Go to Product page](#)

## Datasheet for ABIN7195623 EIF2C3 Protein (His tag)

### Overview

Quantity:	10 µg
Target:	EIF2C3
Origin:	Human
Source:	Baculovirus infected Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This EIF2C3 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human AGO3/Argonaute 3/EIF2C3 Protein (His Tag)
Sequence:	Met 1-Ala 860
Characteristics:	A DNA sequence encoding the human EIF2C3 (Q9H9G7-1) (Met 1-Ala 860) was expressed, with a polyhistidine tag at the N-terminus.
Purity:	> 88 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.

### Target Details

Target:	EIF2C3
Alternative Name:	AGO3/Argonaute 3/EIF2C3 ( <a href="#">EIF2C3 Products</a> )
Background:	Background: Hydroxysteroid sulfotransferase ( SULT2A1 ) is a key enzyme in the testicular and hepatic metabolism of 5alpha-androstenone, which is a major component of the off-odor and off-flavor in pork known as boar taint. Sulfotransferase enzymes catalyze the sulfate

## Target Details

conjugation of many hormones, neurotransmitters, drugs, and xenobiotic compounds. These cytosolic enzymes are different in their tissue distributions and substrate specificities. The gene structure (number and length of exons) is similar among family members. SULT2A1 is a sulfo-conjugating phase II enzyme expressed at very high levels in the liver and intestine, the two major first-pass metabolic tissues, and in the steroidogenic adrenal tissue. SULT2A1 acts preferentially on the hydroxysteroids dehydroepiandrosterone, testosterone/dihydrotestosterone, and pregnenolone and on cholesterol-derived amphipathic sterol bile acids.

Synonym: EIF2C3

Molecular Weight: 99.6 kDa

Pathways: [Fc-epsilon Receptor Signaling Pathway](#), [Regulatory RNA Pathways](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile 20 mM Tris, 500 mM NaCl, pH 7.4, 10 % glycerol

Storage: 4 °C, -20 °C, -80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.