



[Go to Product page](#)

Datasheet for ABIN7195821  
**FLT3 Protein (His tag)**

### Overview

Quantity:	100 µg
Target:	FLT3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This FLT3 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human FLT-3/FLK-2 Protein (His Tag)(Active)
Sequence:	Met 1-Asn 541
Characteristics:	A DNA sequence encoding the extracellular domain of human FLT3 (NP_004110.2) (Met 1-Asn 541) was expressed, with a C-terminal polyhistidine tag.
Purity:	> 85 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to bind biotinylated human FLT3L-His in a functional ELISA.

### Target Details

Target:	FLT3
Alternative Name:	FLT-3/FLK-2 ( <a href="#">FLT3 Products</a> )

## Target Details

---

**Background:** Background: The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD135, also known as FLT-3, FLK-2, is a member of the CD system. CD135 is an important cell surface marker recognized by specific sets of antibodies to identify the types of hematopoietic (blood) progenitors in the bone marrow and its function to differentiate hematopoietic stem cells, which are CD135 negative, from multipotent progenitors, which are CD135 positive. CD135 is a receptor tyrosine kinase type III for the cytokine Flt3 ligand and activates signaling through second messengers by binding to Flt3. Signaling through CD135 is important for lymphocyte development. The encoding gene CD135 is a proto-oncogene to which mutations happened will lead to cancer such as leukemia. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy  
**Synonym:** Receptor-Type Tyrosine-Protein Kinase FLT3; FL Cytokine Receptor; Fetal Liver Kinase-2; FLK-2; Fms-Like Tyrosine Kinase 3; FLT-3; Stem Cell Tyrosine Kinase 1; STK-1; CD135; FLT3; FLK2; STK1

---

**Molecular Weight:** 59.7 kDa

---

**NCBI Accession:** [NP\\_004110](#)

---

**Pathways:** [RTK Signaling](#)

## Application Details

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Lyophilized

---

**Reconstitution:** Please refer to the printed manual for detailed information.

---

**Buffer:** Lyophilized from sterile PBS, pH 7.4

---

**Storage:** 4 °C, -20 °C, -80 °C

---

**Storage Comment:** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

samples are stable at < -20°C for 3 months.