

Datasheet for ABIN7196187

ICOSLG Protein (His tag,Fc Tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	ICOSLG
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This ICOSLG protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Mouse ICOS Ligand/ICOSL Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Lys 279
Characteristics:	A DNA sequence encoding the mouse B7-H2 (NP_056605.1) extracellular domain (Met 1-Lys 279) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized human ICOS at 1 µg/ml (100 µl/well) can bind biotinylated mouse B7-H2 Fc chimera with a linear range of 0.125-1.0 µg/ml.

Target Details


Target:	ICOSLG
---------	--------

Target Details

Alternative Name:	ICOS Ligand/ICOSL (ICOSLG Products)
Background:	<p>Background: Inducible co-stimulator ligand (ICOSL), also known as B7-H2, is a member of the B7 family of co-stimulatory molecules related to B7-1 and B7-2. It is a transmembrane glycoprotein with extracellular IgV and IgC domains, and binds to ICOS on activated T cells, thus delivers a positive costimulatory signal for optimal T cell function. The structural features of ICOSL are crucial for its costimulatory function. Present study shows that ICOSL displays a marked oligomerization potential, resembling more like B7-1 than B7-2. B7-H2-dependent signaling may play an active role in a proliferative response rather than in cytokine and chemokine production. The CD28/B7 and ICOS/B7-H2 pathways are both critical for costimulating T cell immune responses. Deficiency in either pathway results in defective T cell activation, cytokine production and germinal center formation.</p> <p>Immune CheckpointImmune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: WB AntibodiesImmune Checkpoint Targets Co-stimulatory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy Targeted Therapy</p> <p>Synonym: Icosl, AU044799, B7-H2, B7RP-1, B7h, BG071784, GI50, GL50, GL50-B, ICOS-L, KIAA0653, LICOS, Ly115l, Mkiaa0653</p>
Molecular Weight:	54.3 kDa
NCBI Accession:	NP_056605
Pathways:	Production of Molecular Mediator of Immune Response, Activated T Cell Proliferation

Application Details

Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C, -20 °C, -80 °C
Storage Comment:	<p>Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.</p> <p>Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>



Western Blotting

Image 1.