antibodies - online.com







IL1R2 Protein (Fc Tag)





Overview

Quantity:	100 μg
Target:	IL1R2
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This IL1R2 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Mouse IL1R2 Protein (Fc Tag)(Active)	
Sequence:	Met1-Glu355	
Characteristics:	A DNA sequence encoding the mouse IL1R2 (P27931) (Met1-Glu355) was expressed, fused with the Fc region of human IgG1 at the C-terminus.	
Purity:	> 85 % as determined by SDS-PAGE	
Endotoxin Level:	$<$ 1.0 EU per μ g of the protein as determined by the LAL method.	
Biological Activity Comment:	Measured by its ability to bind biotinylated recombinant human IL1α in a functional ELISA.	

Target Details

Target:	IL1R2
Alternative Name:	IL1R2 (IL1R2 Products)

Target Details

Bacl	/~	rol	ind.
וואמ	K (II	() [111(1

Background: Interleukin 1 receptor, type II (IL1R2) also known as CD121b (Cluster of Differentiation 121b) is a cytokine receptor that belongs to the interleukin-1 receptor family. This protein binds interleukin alpha (IL1A), interleukin beta (IL1B), and interleukin 1 receptor, type I (IL1R1/IL1RA), and acts as a decoy receptor that inhibits the activity of its ligands. The pleiotropic cytokine IL1 is produced to regulate development and maintenance of the inflammatory responses, and binds to specific plasma membrane receptors on cells. Two distinct types of IL1 receptors which are able to bind IL1 specifically have been identified, designated as IL1RI (IL1RA) and IL1RII (IL1RB). IL1R1 contributes to IL-1 signaling, whereas the IL-1R2/CD121b has no signaling property and acts as a decoy for IL-1. IL-1R2/CD121b structurally consisting of a ligand binding portion comprised of three Ig-like domains, a single transmembrane region, and a short cytoplasmic domain, is expressed in a variety of cell types including B lymphocytes, neutrophils, monocytes, large granular leukocytes and endothelial cells. Interleukin 4 (IL4) is reported to antagonize the activity of interleukin 1 by inducing the expression and release of this cytokine.

Synonym: CD121b,II1r-2

Molecular Weight:

65.1 kDa

UniProt:

P27931

Pathways:

NF-kappaB Signaling


Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized		
Reconstitution:	Please refer to the printed manual for detailed information.		
Buffer:	Lyophilized from sterile PBS, pH 7.4		
Storage:	4 °C,-20 °C,-80 °C		
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted		
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.		



Western Blotting

Image 1.