
# antibodies .- online.com







## **IL10RB Protein (Fc Tag)**





Go to Product page

$\sim$				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μg
Target:	IL10RB
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL10RB protein is labelled with Fc Tag.

#### **Product Details**

Purpose:	Recombinant Mouse IL10RB/IL10R2 Protein (Fc Tag)	
Sequence:	Met 1-Ser 222	
Characteristics:	A DNA sequence encoding the extracellular domain of mouse IL10RB (NP_032375.2) (Met 1-Ser 222) was fused with the Fc region of human IgG1 at the C-terminus.	
Purity:	> 92 % as determined by SDS-PAGE	
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.	

#### **Target Details**

Target:	IL10RB
Alternative Name:	IL10RB/IL10R2 (IL10RB Products)
Background:	Background: Interleukin 10 receptor, beta subunit (IL10RB/IL-10RB) also known as Cytokine receptor family 2 member 4, Interleukin-10 receptor subunit 2, and cytokine receptor family II,
	member 4, is a subunit for the interleukin-10 receptor. IL10RB/IL-10RB belongs to the cytokine

receptor family. It is an accessory chain essential for the active interleukin 10 receptor complex. Coexpression of this and IL10RA proteins has been shown to be required for IL10-induced signal transduction. Defects in IL10RB/IL-10RB are the cause of inflammatory bowel disease type 25 (IBD25). It is a chronic, relapsing inflammation of the gastrointestinal tract with a complex etiology. It is subdivided into Crohn disease and ulcerative colitis phenotypes. Crohn disease may affect any part of the gastrointestinal tract from the mouth to the anus, but most frequently it involves the terminal ileum and colon. Bowel inflammation is transmural and discontinuous, it may contain granulomas or be associated with intestinal or perianal fistulas. In contrast, in ulcerative colitis, the inflammation is continuous and limited to rectal and colonic mucosal layers, fistulas and granulomas are not observed. Both diseases include extraintestinal inflammation of the skin, eyes, or joints.

Synonym: 6620401D04Rik,AI528744,CRF2-4,Crfb4,D16H21S58,D21S58h,IL-10R2,II10r2

Molecular Weight:

50.5 kDa

NCBI Accession:

NP\_032375


#### **Application Details**

Restrictions:

For Research Use only

#### Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	



### **Western Blotting**

Image 1.