

Datasheet for ABIN7196404  
**IL13RA2 Protein (His tag,Fc Tag)**



[Go to Product page](#)

## Overview

|                               |                                                       |
|-------------------------------|-------------------------------------------------------|
| Quantity:                     | 100 µg                                                |
| Target:                       | IL13RA2                                               |
| Origin:                       | Human                                                 |
| Source:                       | HEK-293 Cells                                         |
| Protein Type:                 | Recombinant                                           |
| Biological Activity:          | Active                                                |
| Purification tag / Conjugate: | This IL13RA2 protein is labelled with His tag,Fc Tag. |

## Product Details

|                              |                                                                                                                                                                                               |
|------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purpose:                     | Recombinant Human IL13RA2/CD213A2 Protein (His & Fc Tag)(Active)                                                                                                                              |
| Sequence:                    | Met 1-Leu 342                                                                                                                                                                                 |
| Characteristics:             | A DNA sequence encoding the extracellular domain of human IL13Rα2 (NP_000631.1) (Met 1-Leu 342) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus. |
| Purity:                      | > 75 % as determined by reducing SDS-PAGE.                                                                                                                                                    |
| Endotoxin Level:             | < 1.0 EU per µg as determined by the LAL method.                                                                                                                                              |
| Biological Activity Comment: | Measured by its binding ability in a functional ELISA. Immobilized recombinant human IL13RA2 at 8 µg/ml (100 µl/well) can bind IL13 with a linear range of 0.25-8.0 ng/ml.                    |

## Target Details

|         |         |
|---------|---------|
| Target: | IL13RA2 |
|---------|---------|

## Target Details

Alternative Name: IL13RA2/CD213A2 ([IL13RA2 Products](#))

Background: Interleukin-13 receptor subunit alpha-2 (IL13RA2/IL-13RA2) is also known as also known as cluster of differentiation 213A2 (CD213A2), IL-13 receptor subunit alpha-2, IL-13R subunit alpha-2, and IL-13RA2. The IL13RA2 is often overexpressed in brain tumors, making IL13ra2 one of the vaccine targets for immunotherapy of glioma. IL13RA2/IL-13RA2 is a cancer-associated receptor that is present in greater than 80% of High Grade Astrocytomas (HGA) and has recently been recognized as a cytokine that predisposes breast cancer cells to metastasize. Expression of IL13Ra2 was rapidly lost from the surface of transduced cells grown in culture. The loss appeared to be related to ligands present in fetal bovine serum in the medium. None of the malignant glioma cell lines cultivated in vitro and tested to date exhibited the IL13Ra2 receptor. A recombinant virus (R5111) enters cells via its interaction with the IL13R  $\alpha$ 2 receptor in a manner that cannot be differentiated from the interaction of wild-type virus with its receptors.

Synonym: CD213A2;CT19;IL-13R;IL13BP;IL13R

Molecular Weight: 65 kDa

NCBI Accession: [NP\\_000631](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile 100 mM Glycine, 10 mM NaCl, 50 mM Tris, pH 7.5

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.