antibodies.com

Datasheet for ABIN7196449 IL18R1 Protein (His tag,Fc Tag)

Image



Overview

Quantity:	100 µg
Target:	IL18R1
Origin:	Mouse
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This IL18R1 protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Mouse IL18R1/CD218a Protein (His & Fc Tag)(Active)		
Sequence:	Met 1-Gly 326		
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Gly 326) of mouse IL18R1 (NP_032391.1) precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.		
Purity:	> 98 % as determined by SDS-PAGE		
Endotoxin Level:	< 1.0 EU per μ g of the protein as determined by the LAL method.		
Biological Activity Comment:	Measured by its ability to bind Cynomolgus IL-18 in a functional ELISA.		

Target Details

IL18R1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7196449 | 09/09/2023 | Copyright antibodies-online. All rights reserved.


Target Details				
Alternative Name:	IL18R1/CD218a (IL18R1 Products)			
Background:	Background: Interleukin-18 receptor 1 (IL18R1) also known as CD218 antigen-like family			
	member A, CDw218a, IL1 receptor-related protein and CD218a, is an interleukin receptor of the			
	immunoglobulin superfamily. IL18R1 is found expressed in lung, leukocytes, spleen, liver,			
	thymus, prostate, small intestine, colon, placenta, and heart, and is absent from brain, skeletal			
	muscle, pancreas, and kidney. High level of expression is found in Hodgkin disease cell lines.			
	This receptor is specifically binds interleukin 18 (IL18), and is essential for IL18 mediated signal			
	transduction. IL18R1 contains 3 Ig-like C2-type (immunoglobulin-like) domains and 1 TIR			
	domain. It is a single-pass type I membrane protein. IFN-alpha and IL12 are reported to induce			
	the expression of this receptor in NK and T cells. The increased expression of IL18R1 may			
	contribute pathogenically to disease and is therefore a potential therapeutic target. The			
	absence of a genetic association in the IL18R1 gene itself suggests regulation from other parts			
	of the genome, or as part of the inflammatory cascade in multiple sclerosis without a prime			
	genetic cause.			
	Synonym: II18ralpha,II1rrp			
Molecular Weight:	63.5 kDa			
NCBI Accession:	NP_032391			
Application Details				
Restrictions:	For Research Use only			
Handling				
Format:	Lyophilized			
Reconstitution:	Please refer to the printed manual for detailed information.			
Buffer:	Lyophilized from sterile PBS, pH 7.4			

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

4 °C,-20 °C,-80 °C

Storage:

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7196449 | 09/09/2023 | Copyright antibodies-online. All rights reserved.



Western	Bl	ott	ing
---------	----	-----	-----

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7196449 | 09/09/2023 | Copyright antibodies-online. All rights reserved.