

Datasheet for ABIN7196497  
**IL21 Receptor Protein (His tag)**



[Go to Product page](#)

## Overview

|                               |  |
|-------------------------------|--|
| Quantity:                     | 100 µg   |
| Target:                       | IL21 Receptor (IL21R)                                |
| Origin:                       | Human  |
| Source:                       | HEK-293 Cells  |
| Protein Type:                 | Recombinant  |
| Biological Activity:          | Active   |
| Purification tag / Conjugate: | This IL21 Receptor protein is labelled with His tag. |

## Product Details

|                              |  |
|------------------------------|--|
| Purpose:                     | Recombinant Human IL-21R Protein (His Tag)(Active)   |
| Sequence:                    | Met 1-Pro 236  |
| Characteristics:             | A DNA sequence encoding the human IL21R (Q9HBE5-1) extracellular domain (Met 1-Pro 236) was expressed, with a polyhistidine tag at the C-terminus.   |
| Purity:                      | > 92 % as determined by reducing SDS-PAGE.   |
| Endotoxin Level:             | < 1.0 EU per µg of the protein as determined by the LAL method.  |
| Biological Activity Comment: | 1. Measured by its ability to bind human IL-21 in a functional ELISA. 2. Measured by its ability to inhibit IL21-induced Interferon gamma secretion by human natural killer lymphoma NK-92 cells. The ED50 for this effect is typically 1-5 ng/ml in the presence of 10 ng/ml of recombinant human IL23. Measured by its ability to inhibit IL21-induced Interferon gamma secretion by human natural killer lymphoma NK-92 cells. The ED50 for this effect is typically 50-200 pg/mL in the presence of 1 ng/ml of recombinant human IL21. |

## Target Details

---

Target: IL21 Receptor (IL21R)

---

Alternative Name: IL-21R ([IL21R Products](#))

---

Background: Interleukin-21 receptor, also known as IL-21 receptor, IL-21R, Novel interleukin receptor, IL21R and NILR, is a single-pass type I membrane protein which belongs to the type I cytokine receptor family and Type 4 subfamily. Interleukin-21 (IL-21) belongs to a family of cytokines that bind to a composite receptor consisting of a private receptor (IL-21R) and the common cytokine receptor gamma chain (  $\gamma(C)$  ). The IL-21R is discovered as a novel member of the class-I-cytokine-receptor family and is selectively expressed in lymphoid tissues. IL-21R shows strong sequence homologies to the interleukin-4 receptor alpha chain gene (IL-4RA). The WSXWS motif of IL-21R appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding. The box 1 motif of IL-21R is required for JAK interaction and / or activation. The IL-21R is widely distributed on lympho-haematopoietic cells and IL21 impacts a number of cell types, including CD8+ memory T cells, NK cells and subsets of CD4 memory T cells. Increased IL21 production is characteristic of certain autoimmune diseases and is likely to contribute to autoantibody production as well as pathological features of autoimmune disease. The critical role of IL21 in promoting humoral immune responses makes it an important focus of potential therapeutic interventions in conditions characterised by overproduction of pathogenic autoantibodies.

Synonym: Interleukin-21 receptor,IL-21 receptor, IL-21R, CD360

---

Molecular Weight: 26.5 kDa

---

Pathways: [JAK-STAT Signaling](#)

---

## Application Details

---

Restrictions: For Research Use only

---

## Handling

---

Format: Lyophilized

---

Reconstitution: Please refer to the printed manual for detailed information.

---

Buffer: Lyophilized from sterile PBS, pH 7.4

---

Storage: 4 °C,-20 °C,-80 °C

---

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.