

Datasheet for ABIN7196972

**MME Protein (His tag)****1** Image[Go to Product page](#)

## Overview

Quantity:	50 µg
Target:	MME
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MME protein is labelled with His tag.

## Product Details

Purpose:	Recombinant Rat CD10/Neprilysin Protein (His Tag)
Sequence:	Tyr52-Trp750
Characteristics:	A DNA sequence encoding the rat MME (P07861) (Tyr52-Trp750) was expressed with a polyhistidine tag at the N-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

## Target Details

Target:	MME
Alternative Name:	CD10/Neprilysin ( <a href="#">MME Products</a> )
Background:	Background: The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are

## Target Details

more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 10 (CD10), also known as Neprilysin and neutral endopeptidase, is a member of the CD system. CD10 is a zinc-dependent metalloprotease enzyme that had function to degrade a number of small secreted peptides such as the amyloid beta peptide. It exist as a membrane-bound protein and have high concentration in kidney and lung tissues. Mutations in the CD10 gene can induce the familial forms of Alzheimer's disease, providing strong evidence for the protein's association with the Alzheimer's disease process. CD10 is also associated with other biochemical processes.

Synonym: MME

Molecular Weight:	82.5 kDa
-------------------	----------

UniProt:	<a href="#">P07861</a>
----------	------------------------

Pathways:	<a href="#">RTK Signaling</a> , <a href="#">Peptide Hormone Metabolism</a> , <a href="#">Regulation of Systemic Arterial Blood Pressure by Hormones</a> , <a href="#">Smooth Muscle Cell Migration</a>
-----------	--

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------

## Handling


Format:	Lyophilized
---------	-------------

Reconstitution:	Please refer to the printed manual for detailed information.
-----------------	--

Buffer:	Lyophilized from sterile PBS, pH 7.4
---------	--------------------------------------

Storage:	4 °C,-20 °C,-80 °C
----------	--------------------

Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
------------------	---



Western Blotting

Image 1.