

Datasheet for ABIN7196999

**TIMP2 Protein (Fc Tag)****1** Image[Go to Product page](#)

## Overview

Quantity:	10 µg
Target:	TIMP2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This TIMP2 protein is labelled with Fc Tag.

## Product Details

Purpose:	Recombinant Human TIMP2/TIMP-2 Protein (Fc Tag)
Sequence:	Cys 27-Pro 220
Characteristics:	A DNA sequence encoding the mature form of human TIMP2 (NP_003246.1) (Cys 27-Pro 220) was expressed with the fused Fc region of human IgG1 at the N-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

## Target Details

Target:	TIMP2
Alternative Name:	TIMP2/TIMP-2 ( <a href="#">TIMP2 Products</a> )
Background:	Background: Tissue inhibitors of metalloproteinases (TIMP) family are natural inhibitors of the matrix metalloproteinases (MMPs), the zinc enzymes involved in extracellular matrix maintenance and remodeling. The TIMP family encompasses four members (TIMP1-4), and

## Target Details

they inhibit most MMPs by forming non-covalent binary complex. TIMP2 is a 22 kDa non N-glycosylated protein expressed by a variety of cell types, and plays a unique role among TIMP family members owing to its functions to regulate cellular responses to growth factors. Findings establish an unexpected, MMP-independent mechanism for TIMP2 inhibition of endothelial cell proliferation in vitro and reveal an important component of the antiangiogenic effect of TIMP2 in vivo. TIMP-2 thus is critical to the maintenance of tissue homeostasis and is involved in the regulation of tumor microenvironment.

Synonym: Metalloproteinase Inhibitor 2, CSC-21K, Tissue Inhibitor of Metalloproteinases 2, CSC-21K, DDC8

Molecular Weight: 48 kDa

NCBI Accession: [NP\\_003246](#)

Pathways: [cAMP Metabolic Process](#)

## Application Details

Restrictions: For Research Use only

## Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile 100 mM Glycine, 10 mM NaCl, 50 mM Tris, pH 7.5

Storage: 4 °C, -20 °C, -80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.