

Datasheet for ABIN7197292 PGC Protein (His tag)



Overview

Quantity:	20 µg
Target:	PGC
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PGC protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human Pepsinogen C/PGC Protein (HEK293 Cells, His Tag)
Sequence:	Met 1-Ala 388
Characteristics:	A DNA sequence encoding the human PGC (NP_002621.1) (Met 1-Ala 388) was fused with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per μ g as determined by the LAL method.

Target Details

Target:	PGC
Alternative Name:	Pepsinogen C/PGC (PGC Products)
Background:	Background: Pepsinogen C, also known as PGC, is an aspartic proteinase that belongs to the peptidase family A1. Pepsinogen C is synthesized in the gastric mucosa as inactive precursors,
	known as zymogens. Pepsinogen C contains a prosegment that serves to stabilize the inactive

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7197292 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

	form and prevent entry of the substrate to the active site. At low PH conditions, Pepsinogen C
	undergoes conversion into active enzyme. Pepsinogen C has been found expressed in all
	regions of the stomach mucosa and also in the proximal duodenal mucosa. In stomach cancer
	tissues and cancer cell lines, the expressions of the pepsinogen genes were decreased or lost,
	in good accordance with their pepsinogen productions. No gross structural changes of the
	pepsinogen genes were observed in these cancers, but the methylation patterns of the
	pepsinogen genes were found to be altered in different ways in different cancers. Serum levels
	of Pepsinogen C are used as a biomarker for certain gastric diseases including Helicobacter
	pylori related gastritis.
	Synonym: PEPC,PGII
Molecular Weight:	42 kDa
NCBI Accession:	NP_002621
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 50 mM Tris, 150 mM NaCl, pH 7.5
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.