

Datasheet for ABIN7197294

PGC Protein (His tag)**1** Image[Go to Product page](#)

Overview

Quantity:	10 µg
Target:	PGC
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PGC protein is labelled with His tag.

Product Details

Purpose:	Recombinant Rat Pepsinogen C/PGC Protein (His Tag)
Sequence:	Met 1-Val 392
Characteristics:	A DNA sequence encoding the rat PGC (NP_579818.1) (Met 1-Val 392) was expressed, fused with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

Target Details

Target:	PGC
Alternative Name:	Pepsinogen C/PGC (PGC Products)
Background:	Background: Pepsinogen C, also known as PGC, is an aspartic proteinase that belongs to the peptidase family A1. Pepsinogen C is synthesized in the gastric mucosa as inactive precursors, known as zymogens. Pepsinogen C contains a prosegment that serves to stabilize the inactive

Target Details

form and prevent entry of the substrate to the active site. At low PH conditions, Pepsinogen C undergoes conversion into active enzyme. Pepsinogen C has been found expressed in all regions of the stomach mucosa and also in the proximal duodenal mucosa. In stomach cancer tissues and cancer cell lines, the expressions of the pepsinogen genes were decreased or lost, in good accordance with their pepsinogen productions. No gross structural changes of the pepsinogen genes were observed in these cancers, but the methylation patterns of the pepsinogen genes were found to be altered in different ways in different cancers. Serum levels of Pepsinogen C are used as a biomarker for certain gastric diseases including Helicobacter pylori related gastritis.

Synonym: PG1,Pg-1,Upg1

Molecular Weight: 42.5 kDa

NCBI Accession: [NP_579818](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile 50 mM Tris, 150 mM NaCl, pH 7.5

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.