

[Go to Product page](#)

Datasheet for ABIN719758

anti-KCNJ9 antibody (AA 61-160) (Biotin)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | KCNJ9 |
| Binding Specificity: | AA 61-160 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This KCNJ9 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human GIRK3 |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Cow, Pig |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | KCNJ9 |
| Alternative Name: | GIRK3 (KCNJ9 Products) |
| Background: | Synonyms: G protein activated inward rectifier potassium channel 3, G protein coupled inward |

Target Details

rectifier potassium channel, GIRK3, Inwardly rectifier K⁺ channel Kir3.3, Inwardly rectifier K⁺ channel KIR3.3, KIR3.3, Potassium channel inwardly rectifying subfamily J member 9, Potassium inwardly rectifying channel subfamily J member 9, Potassium inwardly rectifying channel subfamily J9.

Background: KCNJ9 belongs to the inward rectifier-type potassium channel family and is controlled by G proteins. It associates with another G-protein-activated potassium channel to form a heteromultimeric pore-forming complex. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium, as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium.

Gene ID: 3760

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months