



[Go to Product page](#)

Datasheet for ABIN7198409  
**TNFRSF10A Protein (His tag)**

### Overview

Quantity:	100 µg
Target:	TNFRSF10A
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This TNFRSF10A protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human TRAILR1/TNFRSF10A Protein (His Tag)(Active)
Sequence:	Met 1-Asn 239
Characteristics:	A DNA sequence encoding the human TNFRSF10A (NP_003835.2) extracellular domain (Met 1-Asn 239) was expressed, fused with a polyhistidine tag at the C-terminus.
Purity:	> 92 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to inhibit TRAIL-mediated cytotoxicity using L-929 mouse fibroblast cells treated with TRAIL. The ED50 for this effect is typically 50-200 ng/ml in the presence of 20 ng/ml Recombinant Human TRAIL/TNFSF10.

### Target Details

Target:	TNFRSF10A
---------	-----------

## Target Details

---

Alternative Name:	TRAILR1/TNFRSF10A ( <a href="#">TNFRSF10A Products</a> )
Background:	<p>Background: Tumor necrosis factor receptor superfamily, member 10a (TRAIL R1), also known as TRAIL receptor 1 (TRAIL R1) or CD261 antigen, is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein. TRAIL R1/CD261/TNFRSF10A serves as a receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. TRAIL R1 can promote the activation of NF-kappa-B. TRAIL R1/CD261/TNFRSF10A induces apoptosis of many transformed cell lines but not of normal tissues, even though its death domain-containing receptor, DR4, is expressed on both cell types.</p> <p>Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy</p> <p>Synonym: APO2;CD261;DR4;MGC9365;TNFRSF10A;TRAILR-1;TRAILR1</p>
Molecular Weight:	15.7 kDa
NCBI Accession:	<a href="#">NP_003835</a>
Pathways:	<a href="#">Apoptosis</a> , <a href="#">Positive Regulation of Endopeptidase Activity</a>

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.