antibodies

Datasheet for ABIN7198628 **AKT1 Protein (His tag)**



Overview

Quantity:	50 µg
Target:	AKT1
Origin:	Human
Source:	Baculovirus infected Insect Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This AKT1 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human AKT1/PKB/PKBa Protein (His Tag)(Active)
Sequence:	Met 1-Ala 480
Characteristics:	A DNA sequence encoding the human AKT1 (NP_001014431.1) (Met 1-Ala 480) was fused with a polyhistidine tag at the C-terminus.
Purity:	> 87 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per μ g as determined by the LAL method.
Biological Activity Comment:	1.No Kinase Activity 2. Measured by its ability to bind biotinylated human CD136 in a functional ELISA.
Target Details	

Target: AKT1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7198628 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	AKT1/PKB/PKBalpha (AKT1 Products)
Background:	Background: v-akt murine thymoma viral oncogene homolog 1 (AKT1), or protein kinase B-
	alpha (PKB-ALPHA) is a serine-threonine protein kinase, belonging to the Protein Kinase
	Superfamily. AKT1 is a major mediator of the responses to insulin, insulin-like growth factor 1
	(IGF1), and glucose. AKT1 also plays a key role in the regulation of both muscle cell hypertrophy
	and atrophy. AKT1 activity is required for physiologic cardiac growth in response to IGF1
	stimulation or exercise training. In contrast, AKT1 activity was found to antagonize pathologic
	cardiac growth that occurs in response to endothelin 1 stimulation or pressure overload. AKT1
	selectively promotes physiological cardiac growth while AKT2 selectively promotes insulin-
	stimulated cardiac glucose metabolism. AKT1 deletion prevented tumor initiation as well as
	tumor progression, coincident with decreased Akt signaling in tumor tissues. AKT1 is the
	primary Akt isoform activated by mutant K-ras in lung tumors, and that AKT3 may oppose
	AKT1 in lung tumorigenesis and lung tumor progression. A number of separate studies have
	implicated AKT1 as an inhibitor of breast epithelial cell motility and invasion. AKT1 may have a
	dual role in tumorigenesis, acting not only pro-oncogenically by suppressing apoptosis but also
	anti-oncogenically by suppressing invasion and metastasis.Immune
	Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy
	Synonym: AKT;CWS6;PKB;PKB-ALPHA;PRKBA;RAC;RAC-ALPHA
Molecular Weight:	57 kDa
NCBI Accession:	NP_001014431
Pathways:	PI3K-Akt Signaling, RTK Signaling, TCR Signaling, AMPK Signaling, Interferon-gamma Pathway,
	TLR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin
	Signaling Pathway, Response to Water Deprivation, Regulation of Actin Filament Polymerization
	, Carbohydrate Homeostasis, Glycosaminoglycan Metabolic Process, Cellular Glucan Metabolic
	Process, Regulation of Muscle Cell Differentiation, Cell-Cell Junction Organization, Regulation of
	Cell Size, Skeletal Muscle Fiber Development, Regulation of Carbohydrate Metabolic Process,
	Hepatitis C, Protein targeting to Nucleus, CXCR4-mediated Signaling Events, Signaling Events
	mediated by VEGFR1 and VEGFR2, Negative Regulation of intrinsic apoptotic Signaling,
	Thromboxane A2 Receptor Signaling, Signaling of Hepatocyte Growth Factor Receptor, Positive

Application Details

Restrictions:

For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7198628 | 09/10/2023 | Copyright antibodies-online. All rights reserved. Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 50 mM Tris, 100 mM NaCl, 3 mM DTT, 0.5 mM PMSF, 5 % Glycerol, pH 8.0
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.