

Datasheet for ABIN7198632

AKT3 Protein (AA 106-479) (GST tag,His tag)[Go to Product page](#)**1** Image

Overview

Quantity:	50 µg
Target:	AKT3
Protein Characteristics:	AA 106-479
Origin:	Mouse
Source:	Baculovirus infected Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This AKT3 protein is labelled with GST tag,His tag.

Product Details

Purpose:	Recombinant Mouse AKT3 Protein (aa 106-479, His & GST Tag)
Sequence:	Ala106-Glu479
Characteristics:	A DNA sequence encoding the mouse AKT3 (Q9WUA6-1) (Ala106-Glu479) was expressed with the N-terminal polyhistidine-tagged GST tag at the N-terminus.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method.

Target Details

Target:	AKT3
Alternative Name:	AKT3 (AKT3 Products)
Background:	Background: v-akt murine thymoma viral oncogene homolog 3 (AKT3), also known as PKB-

Target Details

GAMMA, with AKT1/PKBalpha, AKT2/PKBbeta, are the members of Akt kinase family, share extensive structural similarity and perform common as well as unique functions within cells. The Akt signaling cascade initiates at the cell surface when growth factors or other extracellular stimuli activate phosphoinositide 3-kinase (PI3K). AKT3 was discovered to be the predominant isoform activated in sporadic melanomas. Levels of activity increased during melanoma progression with metastatic melanomas having the highest activity. Although mechanisms of AKT3 activation remain to be fully characterized, overexpression of AKT3 and decreased PTEN activity play important roles in this process. Targeted reduction of AKT3 activity decreased survival of melanoma tumor cells leading to inhibition of tumor development, which may be therapeutically effective for shrinking tumors in melanoma patients. AKT2 and AKT3 play an important role in the viability of human malignant glioma cells. Targeting AKT2 and AKT3 may hold promise for the treatment of patients with gliomas.

Synonym: AI851531,D930002M15Rik,Nmf350

Molecular Weight: 71 kDa

Pathways: [PI3K-Akt Signaling](#), [RTK Signaling](#), [TLR Signaling](#), [Hepatitis C](#), [VEGF Signaling](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile 20 mM Tris, 500 mM NaCl, pH 7.4, 10 % glycerol

Storage: 4 °C, -20 °C, -80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.