

Datasheet for ABIN7199568

CD137 Protein (AA 24-86) (His tag)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CD137 (TNFRSF9)
Protein Characteristics:	AA 24-86
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD137 protein is labelled with His tag.

Product Details

Purpose:	Human 4-1BB / TNFRSF9 (24-86) Protein, His Tag
Sequence:	Leu 24 - Cys 86
Characteristics:	Human 4-1BB / TNFRSF9 (24-86), His Tag (41B-H52H9) is expressed from human 293 cells (HEK293). It contains AA Leu 24 - Cys 86 (Accession # Q07011-1).
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	CD137 (TNFRSF9)
Alternative Name:	4-1BB / TNFRSF9 (TNFRSF9 Products)
Background:	Synonyms: TNFRSF9,4-1BB,CD137,CDw137,ILA,

Target Details

Description: 4-1BB is also known as CD137, tumor necrosis factor receptor superfamily member 9 (TNFRSF9), induced by lymphocyte activation (ILA), is a co-stimulatory molecule of the tumor necrosis factor (TNF) receptor superfamily. CD137 can be expressed by activated T cells, but to a larger extent on CD8 than on CD4 T cells. In addition, CD137 expression is found on dendritic cells, follicular dendritic cells, natural killer cells, granulocytes and cells of blood vessel walls at sites of inflammation. The best characterized activity of CD137 is its costimulatory activity for activated T cells. Crosslinking of CD137 enhances T cell proliferation, IL-2 secretion survival and cytolytic activity. Further, it can enhance immune activity to eliminate tumors in mice. CD137 can enhance activation-induced T cell apoptosis when triggered by engagement of the TCR/CD3 complex. In addition, 4-1BB/4-1BBL co-stimulatory pathway has been shown to augment secondary CTL responses to several viruses, and meanwhile augment anti-tumor immunity. 4-1BB thus is a promising candidate for immunotherapy of human cancer. CD137 has been shown to interact with TRAF2.

Molecular Weight: 8.7 kDa

NCBI Accession: [NP_001552](#)

Pathways: [Cancer Immune Checkpoints](#)

Application Details

Application Notes: This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of 8.7 kDa. The protein migrates as 13 kDa under reducing (R) condition due to glycosylation.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Buffer: PBS, pH 7.4

Storage: -20 °C

Storage Comment: -20°C