

Datasheet for ABIN720191  
**anti-RAB7A antibody (AA 101-207)**

3 Images

1 Publication

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	RAB7A
Binding Specificity:	AA 101-207
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAB7A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RAB7
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.

## Target Details

Target:	RAB7A
Alternative Name:	RAB7 ( <a href="#">RAB7A Products</a> )
Background:	Synonyms: RAB7, PRO276, Ras-related protein Rab-7a, RAB7A

## Target Details

---

Background: Key regulator in endo-lysosomal trafficking. Governs early-to-late endosomal maturation, microtubule minus-end as well as plus-end directed endosomal migration and positioning, and endosome-lysosome transport through different protein-protein interaction cascades. Plays a central role, not only in endosomal traffic, but also in many other cellular and physiological events, such as growth-factor-mediated cell signaling, nutrient-transporter mediated nutrient uptake, neurotrophin transport in the axons of neurons and lipid metabolism. Also involved in regulation of some specialized endosomal membrane trafficking, such as maturation of melanosomes, pathogen-induced phagosomes (or vacuoles) and autophagosomes. Plays a role in the maturation and acidification of phagosomes that engulf pathogens, such as S.aureus and M.tuberculosis. Plays a role in the fusion of phagosomes with lysosomes. Plays important roles in microbial pathogen infection and survival, as well as in participating in the life cycle of viruses. Microbial pathogens possess survival strategies governed by RAB7A, sometimes by employing RAB7A function (e.g. Salmonella) and sometimes by excluding RAB7A function (e.g. Mycobacterium). In concert with RAC1, plays a role in regulating the formation of RBs (ruffled borders) in osteoclasts. Controls the endosomal trafficking and neurite outgrowth signaling of NTRK1/TRKA. Regulates the endocytic trafficking of the EGF-EGFR complex by regulating its lysosomal degradation.

Gene ID:	7879
UniProt:	<a href="#">P51149</a>
Pathways:	<a href="#">EGFR Signaling Pathway</a> , <a href="#">Maintenance of Protein Location</a> , <a href="#">SARS-CoV-2 Protein Interactome</a>

## Application Details

---

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 FCM 1:20-100 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

## Publications


Product cited in:	Aboul Naga, Dithmer, Chitadze, Kabelitz, Lucius, Roider, Klettner: "Intracellular pathways following uptake of bevacizumab in RPE cells." in: <b>Experimental eye research</b> , Vol. 131, pp. 29-41, (2015) ( <a href="#">PubMed</a> ).
-------------------	--

## Images




### Immunohistochemistry

**Image 1.** Formalin-fixed and paraffin embedded human colon carcinoma tissue labeled with Rabbit Anti-RAB7 Polyclonal Antibody (ABIN720191) at 1:200 followed by conjugation to the secondary antibody and DAB staining.



#### Immunohistochemistry (Paraffin-embedded Sections)

**Image 2.** Formalin-fixed and paraffin embedded human colon carcinoma tissue labeled with Rabbit Anti-RAB7 Polyclonal Antibody at 1:200 followed by conjugation to the secondary antibody and DAB staining.



#### SDS-PAGE

**Image 3.** Lane 1: mouse liver lysates Lane 2: myeloma cell sp-20 lysates probed with Anti RAB7 Polyclonal Antibody, Unconjugated (ABIN720191) at 1:200 in 4 °C. Followed by conjugation to secondary antibody at 1:3000 90min in 37 °C. Predicted band 23kD. Observed band size: 23kD.