

Datasheet for ABIN720373

anti-UGDH antibody (AA 421-494) (Cy3)



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Alternative Name:

Quantity:	100 μL		
Target:	UGDH		
Binding Specificity:	AA 421-494		
Reactivity:	Human, Mouse, Rat		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This UGDH antibody is conjugated to Cy3		
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence		
	(Paraffin-embedded Sections) (IF (p))		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from human UGDH/UDPGDH		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse, Rat		
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Rabbit		
Purification:	Purified by Protein A.		
Target Details			
Target:	UGDH		

UGDH (UGDH Products)

Target Details

Background:	Synonyms: UDP glucose dehydrogenase, GDH, UDP Glc dehydrogenasev UDP GlcDH, UDP	
	glucose 6 dehydrogenase, UDP-Glc dehydrogenase, UDP-GlcDH, UDP-glucose 6-	
	dehydrogenase, UDPGDH, UGD, Ugdh, UGDH_HUMAN, Uridine diphospho glucose	
	dehydrogenase.	
	Background: The protein encoded by this gene converts UDP-glucose to UDP-glucuronate and	
	thereby participates in the biosynthesis of glycosaminoglycans such as hyaluronan, chondroitin	
	sulfate, and heparan sulfate. These glycosylated compounds are common components of the	
	extracellular matrix and likely play roles in signal transduction, cell migration, and cancer growth	
	and metastasis. The expression of this gene is up-regulated by transforming growth factor beta	
	and down-regulated by hypoxia. Alternative splicing results in multiple transcript	
	variants.[provided by RefSeq, May 2010].	
Gene ID:	7358	
Pathways:	Glycosaminoglycan Metabolic Process	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	